

# **Dark matter: Our method for catching ghostly halos could help unveil what it's made of**

October 22 2020, by Andreea Font

---

Artist's conception of the Milky Way galaxy, which should contain dark matter haloes. Credit: Nick Risinger/NASA

The search for dark matter—an unknown and invisible substance thought to make up the vast majority of matter in the universe—is at a

crossroads. Although it was proposed [nearly 70 years ago](#) and has been searched for intensely—with large particle colliders, detectors deep underground and even instruments in space—it is still nowhere to be found.

But astronomers have promised [to leave "no stone unturned"](#) and have started to cast their net wider out into the galaxy. The idea is to extract information from astrophysical objects that may have witnessed chunks of it as they were passing by. We have just proposed [a new method of doing so](#) by tracing galactic gas—and it may help tell us what it's actually made of.

Physicists believe that dark matter has a propensity to structure itself into a hierarchy of haloes and subhaloes, via gravity. The masses of these clumps fall on a spectrum, with lower mass ones expected to be more numerous. Is there a limit to how light they could be? It depends on the nature of the dark matter particles.

## Warm versus cold

Dark matter cannot be seen directly. We know it exists because we can see the gravitational effects it has on surrounding matter. There are [different theories](#) about what dark matter may actually be. The standard model suggests it is cold, meaning it moves very slowly and only interacts with other matter through the force of gravity. This would be consistent with it being made up of particles known as axions or WIMPS. Another theory, however, suggests it is warm, meaning it moves at higher speeds. One such particle candidate is the [sterile neutrino](#).

If dark matter is cold, a Milky Way-type galaxy could harbour one or two subhaloes weighing as much as  $10^{10}$  Suns, and most likely hundreds with masses of around  $10^8$  Suns. If dark matter is warm, haloes lighter

than around  $10^8$  Suns cannot form easily. So tallying light mass dark haloes can tell us something about the nature of dark matter.

Artist's impression of the expected dark matter distribution around the Milky Way, seen as a blue halo. Credit: ESO/L. Calçada, [CC BY-ND](#)

## **Halo imprints**

We believe that the existence of lower mass haloes can be revealed by carefully planned observations. Astronomers have already got pretty good at this game of hide and seek with dark matter haloes and have devised observations to pick up the damage they leave behind.

To date, observations have targeted mostly the changes in the distribution of stars in the Milky Way. For example, the Large Magellanic Cloud, a smaller galaxy orbiting ours, seems to have a dark matter halo which is massive enough to [trigger an enormous wake](#) – driving the stars from across vast regions to move in unison.

A few of the smaller dark matter haloes thought to be whizzing inside

the Milky Way may occasionally pierce through large stellar features, such as globular clusters (spherical collection of stars), leaving [tell-tale gaps](#) in them. Dark matter haloes can also affect how light bends around astrophysical objects in a process called [gravitational lensing](#).

But the signals left in the stellar distributions are weak and prone to confusion with the stars' own motions. Another way to probe the effect of haloes is by looking at the galactic gas it affects. Galaxies have [plenty of hot gas](#) (with a temperature of around  $10^6$  degrees Kelvin) which extends out to their edge, providing a wide net for catching these dark matter haloes.

A galaxy cluster with dark matter mapped in blue and bright X-rays in pink.

Credit: Smithsonian/wikipedia, [CC BY-SA](#)

Using a combination of analytical calculations and computer simulations, we have shown that dark haloes heavier than  $10^8$  solar masses can compress the hot gas through which they are moving. These will create local spikes in the density of the gas, which can be picked up by X-ray telescopes. These are predicted to be minute, of the order of a few percent, but they will be within the reach of the upcoming [Lynx](#) and [Athena](#) telescopes.

Our models also predict that the spikes in the density of the [cooler galactic gas](#) (with temperature of around  $10^5$  K) will be even more significant. This means that the cooler gas can record the passage of dark matter haloes even more sensitively than the hot gas.

Another promising way of observing the dark-matter-induced fluctuations in the gas is via the photons (light particles) from the cosmic microwave background—the light left over from the Big Bang. This light [scatters off](#) the highly energetic electrons in the hot gas in a way that we can detect, providing a complementary approach to the other studies.

Over the next few years, this new method can be used to test models of dark matter. Regardless of whether dark matter haloes below  $10^8$  solar masses are found in the numbers predicted or not, we will learn something useful. If the numbers match up, the standard cosmological model would have passed an important test. If they are missing, or are far fewer than expected, the standard model would be ruled out and we'll have to find a more viable alternative.

Dark matter remains a mystery, but there's a huge amount of work going

into solving it. Whether the answer will come from instruments on Earth or astrophysical probes, it will no doubt be one of the most important discoveries of the century.

This article is republished from [The Conversation](#) under a Creative Commons license. Read the [original article](#). *This story is part of [Science X Dialog](#), where researchers can report findings from their published research articles. [Visit this page](#) for information about ScienceX Dialog and how to participate.*

Provided by The Conversation

Citation: Dark matter: Our method for catching ghostly halos could help unveil what it's made of (2020, October 22) retrieved 23 April 2024 from <https://sciencex.com/news/2020-10-dark-method-ghostly-halos-unveil.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--