


Western fires are burning higher in the mountains at unprecedented rates in a clear sign of climate change

25 May 2021, by Mojtaba Sadegh, John Abatzoglou and Mohammad Reza Alizadeh



Forest fires advanced to higher elevations as the climate dried from 1984 to 2017. Every 200 meters equals 656 feet. Credit: Mojtaba Sadegh, [CC BY-ND](#)

The Western U.S. appears headed for another [dangerous fire season](#), and a new study shows that even high mountain areas once considered too wet to burn are at increasing risk as the climate warms.

Nearly two-thirds of the U.S. West is in [severe to exceptional drought](#) right now, including large parts of the Rocky Mountains, Cascades and Sierra Nevada. The situation is so severe that the Colorado River basin is on the verge of its [first official water shortage declaration](#), and forecasts suggest another hot, dry summer is on the way.

Warm and dry conditions like these are [a recipe for wildfire disaster](#).


In a [new study](#) published May 24, 2021, our team of [fire and climate scientists and engineers](#) found that forest fires are now reaching higher, normally wetter elevations. And they are burning there at rates unprecedented in recent fire history.

While some people focus on [historical fire suppression](#) and other forest management practices as reasons for the West's worsening fire problem, these high-elevation forests have had little human intervention. The results provide a clear indication that climate change is enabling these normally wet forests to burn.

As wildfires creep higher up mountains, another tenth of the West's forest area is now at risk, according to our study. That creates new hazards for mountain communities, with impacts on downstream water supplies and the plants and wildlife that call these forests home.

Rising fire risk in the high mountains

In the [new study](#), we analyzed records of all fires larger than 1,000 acres (405 hectares) in the mountainous regions of the contiguous Western U.S. between 1984 and 2017.



On average, fires have spread 826 feet (252 meters) higher into the mountains in recent decades, exposing an additional 31,400 square miles (81,500 square kilometers) of forests to fire. Credit: Mojtaba Sadegh, [CC BY-ND](#)

The amount of land that burned increased across all elevations during that period, but the largest increase occurred above 8,200 feet (2,500 meters). To put that elevation into perspective, Denver—the mile-high city—sits at 5,280 feet, and Aspen, Colorado, is at 8,000 feet. These high-elevation areas are largely remote mountains and forests with some small communities and ski areas.

The area burning above 8,200 feet more than tripled in 2001-2017 compared with 1984-2000.

Our results show that climate warming has diminished the high-elevation flammability barrier—the point where forests historically were too wet to burn regularly because the snow normally lingered well into summer and started falling again early in the fall. Fires advanced about 826 feet (252 meters) uphill in the Western mountains over those three decades.

The [Cameron Peak Fire in Colorado](#) in 2020 was the state's largest fire in its history, burning over 208,000 acres (84,200 hectares) and is a prime example of a high-elevation forest fire. The fire burned in forests extending to 12,000 feet (3,650 meters) and reached the upper tree line of the Rocky Mountains.

We found that rising temperatures in the past 34 years have helped to extend the fire territory in the West to an additional 31,470 square miles (81,500 square kilometers) of high-elevation forests. That means a staggering 11% of all Western U.S. forests—an area similar in size to South Carolina—are susceptible to fire now that weren't three decades ago.


Can't blame fire suppression here

In lower-elevation forests, several factors contribute to fire activity, including the presence of more people in wildland areas and a history of fire suppression.

In the early 1900s, Congress commissioned the U.S. Forest Service to [manage forest fires](#), which resulted in a focus on suppressing fires—a policy that continued through the 1970s. This caused flammable underbrush that would normally be cleared out by occasional natural blazes to accumulate. The increase in biomass in many lower elevation forests across the West has been associated with increases in [high-severity fires and megafires](#). At the same time, [climate warming has dried out forests in the Western U.S.](#), making them more prone to large fires.

U.S. wildland area burned each year

More U.S. wildland acres burned in 2015 than any other year on record. The year 2020 was a close second. The majority of these fires are in the West and Alaska.



Prior to 1983, federal wildland fire agencies did not track official wildfire data using current reporting processes. Credit: Chart: The Conversation/CC-BY-ND Source: National Interagency Coordination Center

By focusing on high-elevation fires, in areas with little history of fire suppression, we can more clearly see the influence of climate change.

Most high-elevation forests haven't been subjected to much fire suppression, logging or other human activities, and because trees at these high elevations are in wetter forests, they historically have long [return intervals between fires](#), typically a century or more. Yet they experienced the highest rate of increase in fire activity in the past 34 years. We found that the increase is strongly correlated with the observed warming.

High mountain fires create new problems

High-elevation fires have implications for natural and human systems.

High mountains are natural water towers that normally provide a sustained source of water to millions of people in dry summer months in the Western U.S. The scars that wildfires leave behind—known as burn scars—affect how much snow can accumulate at high elevations. This can influence the timing, [quality and quantity of water](#) that reaches reservoirs and rivers downstream.

High-elevation fires also remove standing trees that act as anchor points that normally stabilize the snowpack, [raising the risk of avalanches](#).

The loss of tree canopy also exposes mountain streams to the Sun, [increasing water temperatures](#) in the cold headwater streams. Increasing stream temperatures can harm fish and the larger wildlife and predators that rely on them.

Climate change is increasing fire risk in many regions across the globe, and [studies show that this trend will continue](#) as the planet warms. The increase in fires in the high mountains is another warning to the U.S. West and elsewhere of the risks ahead as the climate changes.

More information: Mohammad Reza Alizadeh et al, Warming enabled upslope advance in western US forest fires, *Proceedings of the National Academy of Sciences* (2021). [DOI: 10.1073/pnas.2009717118](#)

This article is republished from [The Conversation](#) under a Creative Commons license. Read the [original article](#).

This story is part of [Science X Dialog](#), where researchers can report findings from their published research articles. [Visit this page](#) for information about ScienceX Dialog and how to participate.

Provided by The Conversation

APA citation: Western fires are burning higher in the mountains at unprecedented rates in a clear sign of climate change (2021, May 25) retrieved 3 December 2021 from <https://sciencex.com/news/2021-05-western-higher-mountains-unprecedented-climatechange.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.